



A service of the National Library of Medicine  
and the National Institutes of Health

My NCBI  
[Sign In] [Regis]

[All Databases](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[OMIM](#)[PMC](#)[Journals](#)[Book](#)

Search  for

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)

Display  Show  Sort by  Send to

All: 1 Review: 0

About Entrez  
NCBI Toolbar

Text Version

Entrez PubMed  
Overview  
Help | FAQ  
Tutorials  
New/Noteworthy  
E-Utilities

PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
Special Queries  
LinkOut  
My NCBI

Related Resources  
Order Documents  
NLM Mobile  
NLM Catalog  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

☐ 1: [Med Hypotheses](#). 2005;64(3):458-63.

[Related Articles, Links](#)

**ELSEVIER**  
**FULL-TEXT ARTICLE**

Comment in:

- [Med Hypotheses](#). 2005;65(6):1201-2.

## Add Alzheimer's disease to the list of autoimmune diseases.

### D'Andrea MR.

Johnson and Johnson Pharmaceutical, Research and Development, Drug  
Discovery, Spring House, PA 19477-0776, USA. mdandrea@prdus.jnj.com

A sole pathological event leading to Alzheimer's disease (AD) remains undiscovered in spite of decades of costly research. In fact, it is more probable that the causes of AD are the result of a myriad of intertwining pathologies. However, hope remains that a single awry event could lead to the many pathological events observed in AD brain tissues thereby creating the presentation of simultaneous pathologies. Age-related vascular diseases, which include an impaired blood-brain barrier (BBB), are a common denominator associated with various degrees of dementia, including AD. Recently, a key finding not only demonstrated the anomalous presence of immunoglobulin (Ig) detection in the brain parenchyma of AD tissues but, most importantly, specific neurons that showed degenerative, apoptotic features contained these vascular-derived antibodies. In addition, subsequent studies detected classical complement components, C1q and C5b-9, in these Ig-positive neurons, which also were spatially more associated with reactive microglia over the Ig-negative neurons. Thus, it is possible that the mere presence of anti-neuronal autoantibodies in the serum, whose importance had been previously dismissed, may be without pathological consequence until there is a BBB dysfunction to allow the deleterious effects of these autoantibodies access on their targets. Hence, these observations suggest autoimmunity-induced cell death in AD.

PMID: 15617848 [PubMed - indexed for MEDLINE]

Display  Show  Sort by  Send to